

University of Groningen

Human cell-based in vitro systems for vaccine evaluation

Tapia Calle, María Gabriela

DOI:
[10.33612/diss.100812074](https://doi.org/10.33612/diss.100812074)

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version
Publisher's PDF, also known as Version of record

Publication date:
2019

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):
Tapia Calle, M. G. (2019). *Human cell-based in vitro systems for vaccine evaluation*. [Thesis fully internal (DIV), University of Groningen]. University of Groningen. <https://doi.org/10.33612/diss.100812074>

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.




Chapter 6



Discussion

Discussion

Much of the current knowledge we have about vaccines comes from studies in animal models, mainly murine models. While fundamental immunological concepts derived from studies using inbred, knock out, and transgenic mice have been pivotal to extend our understanding of immune responses, there are multiple reasons to favor the development and use of animal-free alternatives (Box 1).



Box 1. Reasons to look for animal-free alternatives

As of 2010, the European Union (EU) adopted a new legislation concerning the “protection of animals used for scientific purposes” (Directive 2010/63/EU). This was mainly based on the 3Rs principle to replace, reduce, and refine the use of animals for scientific research. This legislation also encourages the development and implementation of alternative methods for animal testing ^[1].

In the context of vaccine development, the use of alternatives to animal models does not mean the complete replacement of animal testing, but rather the inclusion of auxiliary platforms that can aid in the screening of vaccines and vaccine lots. For example, identifying vaccines with low immunogenic potential by employing animal-free alternatives would contribute to the use of fewer animals. Vaccine-related use of animal models is not only restricted to vaccine development but also includes

the monitoring of the production processes of new vaccines and quality control prior to vaccine batch release. This translates into a constant need of animals in order to check the quality of routinely produced vaccine lots (e.g., diphtheria, tetanus, pertussis, Tick-borne encephalitis virus and polio vaccines) [2]. In the context of vaccine development and production, *in vitro* systems can be extremely valuable as they represent an animal-free alternative to assess quality, predict responses and help in elucidating immunological and cellular mechanisms of vaccines and adjuvants.

Given the increased need for alternatives, the ultimate goal of this thesis was to establish a human cell-based system to assess immunological responses to vaccines *in vitro*. We envisioned an approach in which we could dissect different parts of the immune response by focusing on different types of immune cells. Using monocyte-derived dendritic cells (MoDCs) generated from human peripheral mononuclear cells (PBMCs), we first established an *in vitro* platform addressing vaccine-evoked innate immune responses. Subsequently, we developed a platform which allowed us to assess adaptive responses, more specifically T cell-mediated immune responses. We then assessed the performance and suitability of the established platforms by comparing and evaluating immune responses to different types of vaccines. Lastly, we determined how the responses to selected vaccines observed in the human *in vitro* system related to *in vivo* responses as measured in mice. Figure 1 is a graphical summary of the main results obtained from the establishment of our *in vitro* system, also referred as vaccine evaluation system (VES).

Establishing the *in vitro* system

To develop a vaccine evaluation system (VES), we first focused on innate immune responses using antigen presenting cells (APCs). The innate immune system is the first line of defense against pathogens and its cellular component is composed of different cell types (i.e., DC, monocytes, macrophages and natural killer cells) equipped with receptors to sense “non-self” microbial components [3,4]. These cells do not only play a key role in controlling the spread of pathogens; but more importantly create an optimal environment for the generation of antigen presentation with the upregulation of co-stimulatory molecules (i.e., MHCII, CD80, CD86). Moreover, they secrete cytokines and chemokines (i.e., IFN γ , TNF α , IL-12, IL-4, IL-10, CCL-21, CXCL-12, CCR7), that enable the activation and differentiation of adaptive immune players, T and B cells [5,6]. DCs are known to be professional APCs, hence pivotally connect the innate and adaptive responses and crucially determine the magnitude and type of the adaptive immune responses [7–9].



Previously, the use of APCs from cell lines and primary cells has allowed assessing innate immune responses *in vitro*, enabling to understand and model basic immunological responses e.g., the differentiation process of monocytes into APCs [10,11]. The use of APCs has been studied as an alternative to the use of animal testing for vaccine registration and batch release. These studies have primarily focused on bacterial vaccines (i.e., pertussis and *Haemophilus influenzae*) and have favored the use of cell lines over primary cells. The authors claimed that the use of cell lines has the advantage of avoiding safety issues due to unknown infectious pathogens potentially present in blood. Another reason they state is the reproducibility, as opposed to the intrinsic donor-to-donor variation found in primary cells [12,13]. Human primary DCs, on the other hand, have been used to study viral vector vaccines [14–17] and adjuvanted vaccines [12,14,18]. These studies, however, have mainly focused on the mechanism of action of the vector or adjuvant and not on the intrinsic characteristics of the vaccines themselves. Hence, we investigated in **Chapter 2** whether MoDCs could serve as a cellular platform to assess and dissect the immunological properties of vaccines *in vitro* by using two commercially available non-adjuvanted vaccines; namely WIV and SU vaccines.

In establishing this platform, our first choice was to use cell lines as opposed to primary cells, since the first ones are fast and easy to use, cell availability is not a concern and reproducibility is expected to be high. However, even the most promising DC-like cell line, MUTZ-3 [19–25], was found to be unsuitable for our purposes. Therefore, we turned to MoDCs generated from human primary cells. Despite the laborious process to isolate and differentiate cells into DCs, human primary cells have the advantage of retaining many characteristics of the cells *in vivo*. And although using primary DCs brings the possible disadvantage of donor to donor variation, it also gives the opportunity to capture human heterogeneity. In testing the MoDCs, we found that in contrast to the MUTZ-3 cell line, these cells were a suitable platform for screening immunogenic properties of vaccine candidates. MoDCs exposed to WIV and SU vaccines readily displayed vaccine-specific response patterns, reflected in different readouts. WIV could successfully stimulate the expression of genes associated with viral immune responses including MYD88, IRF7, and STAT1, upregulate co-stimulatory molecules on the DC surface like MHCII, CD80 and CD86 and induce the production of cytokines, whilst SU displayed a rather poor capacity to induce such innate immune responses. Notably, the effects of WIV and SU on human DCs *in vitro* were in concordance with those previously reported *in vivo* for mouse DCs [26] and from clinical studies [27,28]. Lastly, we showed that freshly isolated and freeze-thawed PBMCs could be used for MoDCs generation and both performed equally well. This indicates that it is possible to cryopreserve

batches of PBMCs for later use and the repetition of tests, thereby greatly increasing the practical feasibility of a MoDC-based vaccine evaluation system.

The established MoDC-based *in vitro* platform can recapitulate vaccine-specific activation patterns upon stimulation. By allowing to capture the human heterogeneity, this platform represents a useful tool to assess vaccine-induced mechanisms and to estimate *in vivo* responses to novel vaccine formulations.

Having established a platform suitable to dissect innate responses to vaccines, the next logical step was to focus on the adaptive immune




Figure 1. A human cell-based system to assess innate and adaptive immune responses *in vitro*. Focusing on innate responses, we found that; **1.** Human primary monocyte-derived DCs are a more suitable platform to assess the stimulatory properties of vaccines than the MUTZ-3 cell line; **2.** The use of MoDCs enables the discrimination between high and low immunogenic influenza vaccines (WIV and SU) when measuring different parameters (surface markers, gene expression, and cytokine production); **3.** Freshly isolated and frozen/thawed PBMCs are equally suitable for the generation of MoDCs and respond to vaccines in similar ways. To characterize and assess adaptive responses, we established a system making use of long-term cultures of whole PBMCs. Upon stimulation with different vaccines, this system enables the; **4.** Expansion of vaccine-specific T cell responses over time; **5.** Evaluation of the capacity of vaccines to induce activation of and expression of cytotoxicity markers and IFN γ production by T cells; **6.** Phenotyping of responding T cell subsets (naïve, T_{CM}, T_{EM}, and T_{EMRA}); **7.** Evaluation of the induction of T_{FH} responses. Image created with Biorender.

responses, as these develop subsequently after DC stimulation. Different from innate immune responses, which are quickly generated upon encounter of a pathogen but are unspecific [29], adaptive responses develop slowly, yet they are specific and build-up immunological memory [30,31]. The ultimate goal of vaccination is the generation of antigen-specific (memory) immune responses. To achieve this, vaccines mimic the events that take place during infection. Adaptive responses are divided into T cell-mediated and antibody responses [32,33]. For many infectious diseases (i.e., HIV, malaria, tuberculosis, hepatitis and influenza), T cell-mediated responses have shown to be an important requirement for optimal immune protection [34–42]. A T cell-inducing vaccine should ideally be able (among others) to activate and generate cells that can recognize and kill infected cells. However, vaccines should not only induce effector T cells like cytotoxic T lymphocytes (CTLs) and proper memory T cells; but also, induce proper CD4 responses, by eliciting the right type of T helper response (i.e., T_{H1} , T_{H2}). Additionally, vaccines should also provide enough stimulation for the activation of T_{FH} responses. The magnitude and quality of these T_{FH} responses have a direct effect on the generation and maintenance of germinal centers required for proper B cell activation and thus, the generation of high-affinity antibodies [43–46].

Given these facts, in **Chapter 3**, we developed an *in vitro* platform to assess vaccine-induced T cell responses. Traditionally, T cell responses to vaccines have been studied *in vitro* using DC-T cell co-cultures where DCs are generated from (human) PBMCs, pulsed with antigen and later brought in contact with autologous T cells [47–51]. This is, however, a rather tedious and time-consuming procedure. We therefore studied, whether vaccine effects on T cells could also be measured in long-term cultures of unfractionated PBMCs. Next to being simple and straightforward, this approach enables the cross-talk of multiple immune cell types, which might be beneficial for optimal T cell responses.

Using this *in vitro* platform, we observed that influenza antigen-specific T cells present in human PBMCs, when stimulated with suitable vaccines, expanded over time, adopted an activated phenotype and started to produce IFN γ and the cytotoxicity marker CD107. Given the fact that our blood donors had most likely been exposed to influenza before we assumed that the responding cell population would have been memory T cells. Indeed, characterization of the responding T cells revealed that they almost exclusively expressed the memory marker CD45RO with the majority of CD4 T cells being of the effector memory phenotype and the majority of the CD8 T cells being of the central memory phenotype. This result is well in line with responses to influenza vaccines observed *in vivo*

[52–57]. Remarkably, we also detected a response of circulating T_{FH} cells to vaccines in terms of an increase in the frequency of $ICOS^+CD4^+CXCR5^+$ cells and induction of IL-21 production in this cell population. The number and activation status of circulating T_{FH} cells early after vaccination is known to be indicative of later antibody responses [43–46] and the possibility to study vaccine effects on this cell population *in vitro* is thus highly relevant. Using WIV and split influenza vaccine, we observed that the T cells in our *in vitro* system responded in distinct ways to different vaccine formulations bearing the same antigens. Both vaccines were equally potent in activating $CD4^+$ and $CD8^+$ T cells and T_{FH} cells and in inducing $IFN\gamma$ production in $CD4^+$ T cells. However, WIV was superior to split vaccine in activating $IFN\gamma$ production in $CD8^+$ T cells, especially in T_{CM} and T_{EM} . This was well in line with our expectations since WIV is known to be a potent activator of DCs as shown in **Chapter 2** and has the capacity to directly deliver antigen to the cytoplasm from where it can fuel into the MHC class I presentation pathway [58].

Thus, the developed *in vitro* platform for the assessment of T cell responses to vaccines can provide detailed information about vaccine effects on various T cell populations, including T_{FH} cells, which are decisive for antibody responses. As such, the platform can be a valuable tool for assessing vaccine mechanisms *in vitro* and for selecting promising vaccine candidates, at least if the vaccines contain antigens to which blood donors likely have memory T cell responses.

Applying and validating the *in vitro* system

As reported from preclinical and clinical trials, influenza vaccines derived from different virus subtypes differ in immunogenicity [23,42–45]. Virus strain-related differences in immunogenicity have important consequences for vaccine formulations as they might necessitate adjusting the amount of antigen or the addition of an adjuvant to achieve adequate protection [59–65]. Previous work, reports intrinsic features of the respective vaccines to account for the observed strain-specific differences in immunogenicity [66]. These studies however, did not employ vaccines which had been produced in a consistent way; hence, an adequate comparison could not be performed and a proper elucidation of the immunological differences between influenza virus subtypes remained to be performed. In **Chapter 4**, we set out to exploit the potential of our previously established *in vitro* system to evaluate and compare vaccines derived from different influenza virus subtypes.

For this purpose, we performed a head-to-head comparison of WIV and SU vaccines derived from H1N1pdm09, H3N2, H5N1 and H7N9 influenza virus

subtypes produced under standardized conditions. Using a systematic approach, we first focused on the physicochemical characteristics of the vaccines; subsequently, we evaluated their immunological properties using *in vitro* and *in vivo* approaches. Concerning WIV, our comparisons showed clear-cut physicochemical and immunological differences that enabled the discrimination of high (H5N1), intermediate (H1N1pdm09, H3N2) and low (H7N9) immunogenic influenza virus subtypes. Yet, differences concerning SU vaccines were less noticeable. By comparing the immunological effects of the different virus subtype vaccines *in vitro* and *in vivo*, we further assessed in how far the established *in vitro* platforms (**Chapter 2 and 3**) could recapitulate the *in vivo* responses. Notably, results of both *in vitro* and *in vivo* readouts correlated well.

In conclusion, through this head-to-head comparison, we gained valuable insights into the intrinsic differences between vaccines derived from different influenza virus subtypes. This comparison will help improve poorly immunogenic influenza vaccines as future experiments can now focus on the use of alternative vaccine modalities, e.g. adjuvants that can improve immunogenicity. We additionally corroborated the suitability of our *in vitro* system as promising animal free-alternatives for vaccine screening and evaluation.

Lastly, **Chapter 5** describes the use of the T cell *in vitro* platform to assess whether vaccine immunogenicity can be improved by using particulate delivery systems and manipulating the particle size. As shown by others^[67,68] and confirmed by us in **Chapter 2**, subunit vaccines, consisting of soluble proteins, are poorly immunogenic. The use of adjuvants can improve immunogenicity. Adjuvants are classified into immunopotentiators and particulate delivery systems; both assist in inducing and modifying immune responses^[69–71]. Particulate antigen delivery systems consist of particles ranging from nano- to micrometer size that can accomplish the delivery and presentation of bound or encapsulated antigens to APCs^[72]. Interestingly, different studies have shown the possibility to steer the type of immune response (T_{H1} , T_{H2}) by manipulating the size, shape or rigidity of particulate delivery systems^[73–76]. Here, we investigated in how far manipulation of size affects antigen uptake by human moDCs and stimulation of human T cells, the latter using the previously established T cell platform.

For this purpose, we used two different vaccines, influenza SU and Hepatitis B (HBsAg), to assess the effect of coupling vaccines to nano- or micro-polystyrene particles (0,5 and 3 μm) on the immune response. Microscopical observations depicted that nano- and microparticle-coupled vaccines, as well as unconjugated vaccines, could be taken up by DCs



with an apparent similar efficiency. Nevertheless, employing our *in vitro* T cell evaluation platform, we observed that influenza vaccine coupled to 3 μm particles was somewhat more immunogenic than unconjugated SU influenza vaccine. This was reflected in an increased number of CD8⁺ T cells producing cytokines and in the total amount of cytokines produced. In contrast, coupling of HBsAg to beads did not affect the magnitude of the T cell responses, except that the frequencies of IFN γ - and IL10-producing T cells were slightly higher when the antigen was coupled to 3 μm beads. All our blood donors most likely had been exposed to influenza several times and thus their PBMCs contained memory T cells which could rapidly respond to encounter of influenza vaccine *in vitro*. In contrast, the donors were most likely naïve for Hepatitis B. Nevertheless, we could still detect T cell responses to HBsAg in some of our donors, albeit at low frequency. This indicates the versatility of the developed *in vitro* platform to assess not only vaccine effects on memory T cells but also the capacity of vaccines to activate naïve T cells. Yet, the experimental setup did not allow us to discriminate whether the observed responses were mediated by naïve or memory T cells. Thus, further experiments are still needed to corroborate the induction of naïve antigen-specific T cell responses upon *in vitro* stimulation with HBsAg and other de novo antigens.

Taken together, size modification of influenza and Hepatitis B vaccines displayed a measurable but rather small effect on the induction of T cell responses. Although we did find a significant effect of coupling of influenza SU to 3 μm beads on the amounts of certain cytokines produced by the stimulated T cells, the biological relevance of this is questionable since the magnitude of the effect was small. Our results might also indicate that particle size is of minor importance for downstream immune responses in human PBMCs.

Gaps, challenges and opportunities

Gaps

In the context of research and development of vaccines, there are three key needs; 1) antigen selection and vaccine design; 2) novel technologies and routes of administrations; 3) clinical studies and data interpretation [77]. In this context, the use and further development of the *in vitro* vaccine evaluation platform presented in this thesis would be an exciting approach to help solving one of these needs: antigen selection and vaccine design.

Unraveling the mechanisms of infection and host-pathogen interactions is paramount for a rational vaccine design. Many infectious diseases for which we do not have efficient vaccines yet, display very complex dynamics that hamper the design of long-term protective vaccines. Examples of this



include pathogens like influenza, dengue, tuberculosis and malaria that exhibit complex pathogenesis, wide-ranging variability and have evolved different strategies to evade the immune system ^[78–80]. Finding a vaccine for these pathogens would require to test a significant pool of antigen targets in order to find protective responses. Hence, a rational vaccine approach should be based on a clear understanding of the immunogenicity of potential key antigens and their interaction with the host. Recognizing the human innate and adaptive responses to specific antigens is thus key to select and design effective vaccines.

In this context, VES, which focusses on innate (DCs) and adaptive (T cells) responses could potentially serve three purposes;

Select vaccines

By comparing different vaccine formulations using cells of the same donors, VES allows evaluating whether one vaccine formulation is better than another one. It allows the comparison of vaccines more easily than in animal experiments and clinical trials. Furthermore, it permits the evaluation of vaccine candidates in donors from different contexts (i.e., sex, age, ethnicity, health status). For example, it would facilitate testing different vaccine formulations in particular target groups, e.g. the elderly, which would be challenging to do by means of clinical trials.

Reveal vaccine mechanisms and ways to achieve the needed protective responses

Using VES we can address the characteristics and functions of potent immunogens and evaluate approaches to steer towards specific immune responses. On the one hand, we can reveal the mechanistic nature of the protective immune response by dissecting innate immune pathways (including involved PRR, surface markers, cytokine, chemokines) activated and/or induced by different types of vaccines. Moreover, we can evaluate the possibility of steering the phenotype (i.e., T_{H1} , T_{H2}) of an existing immune response to favor efficient protection. For instance, by assessing earlier identified pathways related to the activation of the innate immune system in PBMCs of young vs. elderly individuals, we could reveal mechanistic differences between the cell patterns of these two target groups.

Assess vaccine quality

For many vaccines, the use of animal models to verify vaccine batch quality is still mandatory. Surprisingly, the number of animals required for these purposes outnumbers that used for scientific research ^[81]. Currently, there are several initiatives to make such experiments

oblivious by following a 'consistency approach' ^[82-84], this means the "in-process" monitoring during vaccine production rather than focusing on final batch testing. This monitoring relies solely on *in vitro* biochemical, physicochemical and cellular tests to guarantee the consistency of the new batch with previous ones ^[83-85]. An *in vitro* approach, as described in this thesis, would allow determining the capacities of vaccine batches to stimulate APCs or T cells and would as such contribute to a reduction of animal use.

Challenges

It is clear that our *in vitro* platform is still a reductionist approach to capture the complexity of the human immune system; and that it cannot fully recapitulate complex interactions. Essential components that can affect the immune response might not be captured (i.e., lymph nodes, germinal centers, gut microbiota). Furthermore, additional validation studies need to be performed to determine in how far the *in vitro* results reflect *in vivo* responses and to what degree they are predictive for vaccination outcome. For instance, in our *in vitro* studies we have shown that WIV was superior to SU vaccine in stimulating APCs and T cells and these results are well in line with *in vivo* results from previous preclinical and clinical studies ^[27,59,86,87]. However, what is still missing is a head-to-head comparison of immune responses *in vitro*, *ex vivo* and *in vivo* to a specific vaccine. For this, *in vitro* effects of vaccines on PBMCs from unvaccinated individuals should be compared to effects of the vaccine on PBMCs measured *ex vivo* shortly after vaccination and be related to the final vaccination outcome in terms of antibody titers and/or number of T cells induced. With this, we would have proof that a vaccine that performs well *in vitro* does the same *in vivo*. Additionally, it would be required to define a response profile having predictive value for the *in vivo* situation. For this purpose, we could potentially make use of the biomarkers recently discovered by systems vaccinology ^[88].

Opportunities

The recent boom of systems vaccinology has expanded our knowledge on the molecular basis of the response to conventional commercial vaccines for influenza ^[89-92] and yellow fever ^[93,94] and vaccines under development against HIV, dengue and Ebola ^[95-97]. By using high-throughput 'omics' approaches on PBMCs from vaccinated individuals, systems vaccinology has identified key biomarkers and responses predictive of vaccine-induced protection. It is now clear that PBMCs are an important source of information; and that early immune signatures can be used to predict later responses (i.e., protective levels of antibodies) ^[89-94]. The most relevant examples are the identification of GCN2 and CaMKIV as predictors of vaccine responses.



Expression levels of GCN2 in PBMCs from recently vaccinated individuals were shown to be predictive of CD8 T cells responses and neutralizing antibody titers against yellow fever vaccines [93,94,98]. CaMKIV expression levels have been shown to inversely correlate with antibody responses in individuals vaccinated with influenza trivalent inactivated vaccines (TIV) and could reliably predict vaccination outcome as early as three days after vaccination [99]. More recently, the generation and activation of circulating T_{FH} (identified as $CD4^+$, $CXCR5^+$, $ICOS^+$, $IL-21^+$) cells shortly after vaccination were found to correlate with the quality and magnitude of the resulting antibody responses in the context of influenza [46,100–102] and HIV vaccines [103,104].

High throughput “omics” approaches are hypothesis-generating; this means, they perform analysis based on significant amounts of acquired data to delineate hypotheses. However, these hypotheses need to be confirmed, dissected and exploited in conventional assays like the *in vitro* system we have developed. For example, evaluating previously identified biomarkers *in vitro* could also help us in understanding their mechanisms of action by correlating them to the generation of new specific cells subsets, their frequency (percentages), activation state and function (cytokines and proliferation potential).

An additional opportunity for VES is to test vaccine effectiveness on an individual level. Vaccine failure and/or vaccine effectiveness can be influenced by many characteristics of the recipient such as age, vaccine history, health status, ethnicity, sex, or pregnancy [105–107]. For instance, infants (below 1 year) do not respond appropriately to meningococcal vaccine; hence, multiple doses are required [108,109]. Also, it is known that with age, the immune system loses its capacity to respond to infections and vaccinations adequately [110]. VES could be employed using PBMC donors from infants or the elderly to evaluate the reasons behind this and to find alternatives (adjuvants) to improve vaccination outcome.

As of today, we have 26 vaccines against different types of human infectious diseases, and 24 additional vaccines are under development (Table 1) [111]. Available vaccines could allow us to further explore our VES, i.e., by checking whether previously described biomarkers (i.e., GCN2 and CaMKIV) also show up in the developed *in vitro* system. Assessing the available vaccines would also be an excellent chance to gain more knowledge into successful mechanisms of protection. Having a clear understanding of how these vaccines work can provide insights into the development of better candidates or the improvement of others. Knowing that these biomarkers correlate well in our system, we could 1) better understand and dissect the induction of vaccine-related responses, 2) check the induction of responses in individuals with different age, sex and geographic context

Table 1. Vaccines available and in the pipeline.

Available Vaccines			Pipeline Vaccines		
Cholera	Haemophilus influenzae type b	Meningococcal meningitis	Campylovacter jejuni	HIV-1	Respiratory Syncytial Virus
Dengue	Human papillomavirus (HPV)	Malaria	Chagas disease	Human Hookworm Disease	Schistosomiasis Disease
Diphtheria	Influenza	Measles	Chikungunya	Leishmaniasis Disease	Shigella
Hepatitis (A, B, E)	Japanese encephalitis	Mumps	Dengue	Malaria	Staphylococcus aureus
Pertussis	Pneumococcal disease	Poliomyelitis	Enterotoxigenic E. coli	Nipah virus	Streptococcus pneumoniae
Rabies	Rotavirus	Rubella	Enterovirus 71	Nontyphoidal Salmonella Disease	Streptococcus pyogenes
Tetanus	Tick-borne encephalitis	Tuberculosis	Group B Streptococcus	Norovirus	Tuberculosis
Typhoid	Varicella	Yellow fever	Herpes Simplex Virus	Paratyphoid fever	Universal Influenza Vaccine

and 3) compare between vaccines, if multiple candidates are available for the same pathogen.

Future Perspectives

Comparing vaccine-related responses in young and old individuals

Vaccines still perform suboptimal in the elderly because general vaccine design and formulation is not aimed at the elderly and the molecular mechanisms behind decreased vaccine responses in elderly remain incompletely understood.

Poor vaccine responses are mainly due to immunosenescence, which describes the fact that all of the components of the immune system suffer from aging and progressive loss of responsiveness^[112,113]. This process is not only reflected in a restricted ability of the aging immune system to protect the body against pathogens but also in a limited response to vaccination^[102,114]. It is known that aging is associated with alterations in the differentiation potential towards memory subsets of T and B cells^[56,115]. The number of naïve T cells decreases together with the TCR repertoire, constriction of this repertoire contributes to a failure to respond to different pathogens^[116]. Regrettably, innate immune cell types also display age-related alternations^[117]. For instance, the function of neutrophils, granulocytes and natural killer (NK) cells is impaired in older adults^[117]. In

addition, in macrophages and DCs, Toll-like receptor (TLR) dysregulation together with lower expression of co-stimulatory molecules have also been demonstrated [118–120].

Due to an impaired immune system, the severity of infectious diseases in the elderly is much higher than in younger adults. Moreover, infections are associated with long-term repercussions affecting daily activities, onset of frailty and decrease of independence [121–124]. Thus, prevention of infectious diseases through vaccination is an important challenge to ensure healthy aging. Yet, traditional vaccines against influenza, *Streptococcus pneumoniae*, herpes zoster, tetanus, diphtheria and pertussis have all been found to be less effective in the elderly [125–128]. In the context of influenza vaccines, a meta-analysis showed that vaccine effectiveness in the elderly (65 + population) is 49% for laboratory-confirmed influenza cases [129] compared to 59% in adults between 18–65 years of age [130]; which indicates the need for more potent vaccines [114,130,131].

The established *in vitro* platforms would be ideal to compare vaccine-related immune responses from young and old individuals and thus give an unprecedented chance to better understand why vaccines perform poorly in the elderly; which is not possible using animal models. This comparison could give us insights into; 1) the mechanistical background of immune-related differences between young and old individuals in their response to vaccine and 2) the evaluation of new strategies to ameliorate the responses to vaccines through new adjuvant compounds. As an additional feature this approach would also allow us to better understand variation within heterogeneous populations with distinct genetic and environmental background. In particular, the different exposure history of young and elderly, including the virus strain they encountered first in life, is expected to have effect on the vaccine-induced responses [132]. Overall, this system would allow to compare different vaccines in a much easier and feasible way than testing them in animal models or in clinical trials.

Expanding towards B cell responses

A remaining challenge is the establishment of an *in vitro* system which allows the evaluation of vaccine-induced B cell responses. Typically, vaccines aim at avidly stimulating B cells to induce the production of protective antibodies [133]; however, B cells can also amplify and suppress immune responses by mechanisms different from antibodies [134–137]. Similar to DCs, B cells possess TLRs, present antigens and produce cytokines, thus exerting regulatory and effector functions [134–138]. Given these features, it would be very interesting to design a B cell-based screening system to assess the immunogenicity of vaccines *in vitro*. Such a platform would allow for zooming in into different aspects of the immune response like

the differentiation of different B cell subsets and their proliferation, the induction of cytokine production, and finally the production of antibodies (Table 2). By combining such a B cell system with the previously established platforms, it would be possible to have an integrated and comprehensive view of key immune cell players in the context of vaccine-induced immune responses.

Table 2. Potential processes to study *in vitro* during B cell-induced responses.

Event	Parameter to measure	Technique	Reference
Naïve B cell activation	MHCII, CD86	Immunofluorescence – Flow cytometry	[139]
Memory B cell activation	MHCII, CD86	Immunofluorescence - Flow cytometry	[140–143]
B cell differentiation	AID, Blimp-1, BAFF, Plasma cells, Plasmablasts	qPCR - Flow cytometry	[139,144–146]
Proliferation	CFSE, Ki-67	DNA synthesis/HTS - Flow cytometry	[139,141,147]
Cytokine production	IL-4, IL-6, IL-10, IL-12, IFN γ , TNF α	ELISpot – ELISA	[142,143]
Antibody production	Antigen-specific IgG, IgM	ELISpot – ELISA	[91,148–150]

Building such a system would require first to confirm whether B cells *in vitro* would respond to vaccines at all using the parameters listed in Table 2. Subsequently, the identification of key downstream mechanisms modulated by vaccines, such as subset differentiation and cytokine production, could need to be performed. Few studies have exploited the potential of B cells *in vitro* ^[139,143,147,151–155], these studies used TLR agonists and not antigens, hence showing the feasibility of a B cell platform with an antigen-independent B cell stimulation approach. Lanzavecchia and colleagues shed light onto important mechanisms of naïve B cell stimulation *in vitro* ^[139,147,151,152]. Mimicking T cell help with CD40L together with TLR9 agonists they showed the possibility to induce B cell activation and thus proliferation and induction of cytokines ^[139]. Others demonstrated that also memory B cells can be activated with a combination of CpG and CD40, and can be differentiated into plasma cells and plasmablasts ^[143,153]. Jung and colleagues went further into the effects of CpG on the different B cell subtypes *in vitro* concluding that this TLR9 agonist had an effect in the terminal differentiation, indicated by an increase in generation and proliferation of plasma cells from naïve and memory B cells. ^[154] As for antibody production, *in vitro* stimulation with a TLR7 ligand induced the secretion of IgM and IgG by naïve B cells ^[155] and IgM, IgG and IgA by memory B cells ^[143].

Antigen-dependent B cell activation *in vitro* has remained challenging until recently. This has been mainly due to haplotype variation in MHCIIIs requiring specific T cells for B cell activation ^[156]. In an effort to generate therapeutic antibodies, a recent study showed the possibility to overcome the stringent requirements for antigen-specific stimulation by using streptavidin nanoparticles conjugated to both CpG and antigen ^[157,158]. Using this approach, the Batistas' lab showed the possibility to identify, quantify and characterize B cells producing antigen specific antibodies. By stimulating memory B cells from healthy donors with CpG in combination with tetanus toxoid or influenza hemagglutinin, they demonstrated the induction of plasma cells able to secrete antigen specific antibodies ^[158].

In an exploratory approach, we have performed experiments on vaccine-stimulated B cells *in vitro*. Our initial results showed the upregulation of different genes related to B cell fate (encoding for Blimp-1, Pax5 and AID) upon stimulation with WIV but not SU vaccines, as well as the induction of proliferation of WIV-treated but not SU-treated cells as measured by CFSE staining. Yet, according to our experience, although possible, the establishment of an *in vitro* B cell system for vaccine evaluation will be a challenging mission. Different from the T cell approach, cell availability is a problem (there are only around 10% of B cells in PBMCs); next to this, there are no studies so far that assess the effect of mitogens nor antigens in entire B cell populations *in vitro*, rather the available papers start from certain subpopulations like naïve or memory B cells ^[139,151,158,159]. This could mean that whole PBMCs are not suitable to assess B cell responses and subpopulations would need to be isolated necessitating the use of several antibodies for the isolation of the required populations and further decreasing the number of cells that can be obtained from a given blood sample.

Optimizing this system will require to carefully phenotype and pinpoint those B cell subsets that could give the most relevant information in the context of vaccine induced responses. Moreover, since antigen-activated B cells may undergo different differentiation fates (i.e. memory B cells, short-lived and long-lived plasma cells) and with different requirements for stimulation (with or without T cell help) ^[101,160,161], it will be necessary to carefully evaluate each of these possibilities to find the best approach to assess B cell vaccine-induced responses *in vitro*.

Concluding remarks

PBMCs can give important information on the immune system and the mechanisms of protection induced by vaccines ^[89,162–166]. In the context of vaccinated individuals, PBMCs have enabled to pinpoint biomarkers and predictive responses for vaccine protection ^[43,44,46,102,167–169].

Given this fact, we hypothesized that PBMCs could represent a strong tool in vaccine development. Using influenza as our main model antigen we therefore developed a MoDC-based system to study vaccine effects on APCs and a system using unfractionated PBMCs to study vaccine effects on T cells. We then used the established platforms together with various vaccine formulations to (1) assess vaccine-induced innate immune cell activation of MoDCs; (2) characterize details of vaccine-induced T cell responses; and (3) validated the system against mice *in vivo* responses. In doing so, we found that human primary cells are a better platform to assess responses to vaccines *in vitro* than cell lines and that activation properties of primary MoDCs distinguished different vaccines formulations. After stimulation of whole PBMCs with different vaccine formulations, T lymphocytes displayed distinct magnitudes of response with respect to proliferation and cytokine production. This system also enabled the evaluation of T_{FH} responses, indicating a potential predictive role for antibody responses. In exploiting this platform, we compared vaccines derived from different influenza virus subtypes revealing that different virus subtypes induce different levels of immunogenicity *in vitro* which reflect their immunogenicity in immunized mice.

Our human cell-based platform to assess responses to vaccines *in vitro*, provides an interesting animal free-alternative to circumvent ethical issues and to fulfill the need of representative models of the human situation (given the poor predictive value of animal models [170–172]). The developed platforms intend to be an auxiliary tool for the screening of vaccine candidates. Identifying poorly immunogenic vaccine candidates *in vitro*, would allow to reduce the number of animals needed for preclinical stages decisively. Moreover, it will enable the selection of strong vaccine candidates with better odds to perform well in a clinical setting. Furthermore, the VES can give important insights into the molecular and cellular mechanisms behind vaccines and adjuvants. Seemingly, an extension of this *in vitro* platform could assist in assessing vaccine responses in risk groups (i.e. elderly), ultimately allowing the development of improved vaccines for specific target populations.

In conclusion, this human cell-based platform to assess responses to vaccines *in vitro*, represents a fast and sustainable system with the ability to reflect/recapitulate responses of the human immune system; hence, its use could potentially aid in solving current issues of vaccine development.

References

- [1] The European parliament and the Council of the European Union. Legislation for the protection of animals used for scientific purposes. 2010.
- [2] Council of Europe. European Pharmacopoeia (Ph. Eur.) 9th Edition | EDQM. 9th ed. n.d.
- [3] Iwasaki A, Medzhitov R. Regulation of Adaptive Immunity by the Innate Immune System. *Science* (80-) 2010;327:291–5. doi:10.1126/science.1183021.
- [4] Wu J, Chen ZJ. Innate Immune Sensing and Signaling of Cytosolic Nucleic Acids. *Annu Rev Immunol* 2014;32:461–88. doi:10.1146/annurev-immunol-032713-120156.
- [5] Morel PA, Turner MS. Designing the optimal vaccine: the importance of cytokines and dendritic cells. *Open Vaccine J* 2010;3:7–17. doi:10.2174/1875035401003010007.
- [6] Curtsinger JM, Mescher MF. Inflammatory cytokines as a third signal for T cell activation. *Curr Opin Immunol* 2010;22:333–40. doi:10.1016/j.coi.2010.02.013.
- [7] Iwasaki A, Medzhitov R. Toll-like receptor control of the adaptive immune responses. *Nat Immunol* 2004;5:987–95. doi:10.1038/nri112.
- [8] Alloatti A, Kotsias F, Magalhaes JG, Amigorena S. Dendritic cell maturation and cross-presentation: timing matters! *Immunol Rev* 2016;272:97–108. doi:10.1111/imr.12432.
- [9] Napolitani G, Rinaldi A, Berton F, Sallusto F, Lanzavecchia A. Selected Toll-like receptor agonist combinations synergistically trigger a T helper type 1-polarizing program in dendritic cells. *Nat Immunol* 2005;6:769–76. doi:10.1038/ni1223.
- [10] Vandebruel R, Hoefnagel MMN. Dendritic cell-based *in vitro* assays for vaccine immunogenicity. *Hum Vaccin Immunother* 2012;8:1323–5. doi:10.4161/hv.21350.
- [11] Hou W, Gibbs JS, Lu X, Brooke CB, Roy D, Modlin RL, et al. Viral infection triggers rapid differentiation of human blood monocytes into dendritic cells. *Blood* 2012;119:3128–31. doi:10.1182/blood-2011-09-379479.
- [12] Hoefnagel MHN, Vermeulen JP, Scheper RJ, Vandebruel RJ. Response of MUTZ-3 dendritic cells to the different components of the *Haemophilus influenzae* type B conjugate vaccine: Towards an *in vitro* assay for vaccine immunogenicity. *Vaccine* 2011;29:5114–21. doi:10.1016/j.vaccine.2011.05.050.
- [13] Hoonakker ME, Verhagen LM, Hendriksen CFM, van Els CACM, Vandebruel RJ, Sloots A, et al. *In vitro* Innate Immune Cell Based Models To Assess Whole Cell Bordetella Pertussis Vaccine Quality: A Proof Of Principle. *Biologicals* 2015;43:100–9. doi:10.1016/j.biologicals.2014.12.002.
- [14] Kastenmuller W, Drexler I, Ludwig H, Erfle V, Peschel C, Bernhard H, et al. Infection of human dendritic cells with recombinant vaccinia virus MVA reveals general persistence of viral early transcription but distinct maturation-dependent cytopathogenicity. *Virology* 2006;350:276–88. doi:10.1016/j.virol.2006.02.039.
- [15] Guerra S, Nájera JL, González JM, López-Fernández LA, Climent N, Gatell JM, et al. Distinct gene expression profiling after infection of immature human monocyte-derived dendritic cells by the attenuated poxvirus vectors MVA and NYVAC. *J Virol* 2007;81:8707–21. doi:10.1128/JVI.00444-07.
- [16] Gruber A, Kan-Mitchell J, Kuhen KL, Mukai T, Wong-Staal F. Dendritic cells transduced by multiply deleted HIV-1 vectors exhibit normal phenotypes and functions and elicit an HIV-specific cytotoxic T-lymphocyte response *in vitro*. *Blood* 2000;96:1327–33.
- [17] Gardner JP, Frolov I, Perri S, Ji Y, MacKichan ML, zur Megede J, et al. Infection of Human Dendritic Cells by a Sindbis Virus Replicon Vector Is Determined by a Single Amino Acid Substitution in the E2 Glycoprotein. *J Virol* 2000;74:11849–57. doi:10.1128/JVI.74.24.11849-11857.2000.
- [18] Hunsawong T, Sunintaboon P, Warit S, Thaisomboonsuk B, Jarman RG, Yoon I-K, et al. Immunogenic Properties of a BCG Adjuvanted Chitosan Nanoparticle-Based Dengue Vaccine in Human Dendritic Cells. *PLoS Negl Trop Dis* 2015;9:e0003958. doi:10.1371/journal.pntd.0003958.
- [19] Larsson K, Lindstedt M, Borrebaeck CA. Functional and transcriptional profiling of MUTZ-3, a myeloid cell line acting as a model for dendritic cells. *Immunology* 2006;117:156–66.
- [20] van Helden SFG, van Leeuwen FN, Figdor CG. Human and murine model cell lines for dendritic cell biology evaluated. *Immunol Lett* 2008;117:191–7. doi:10.1016/j.imlet.2008.02.003.



- [21] Santegoets SJAM, van den Eertwegh AJM, van de Loosdrecht AA, Scheper RJ, de Gruijl TD. Human dendritic cell line models for DC differentiation and clinical DC vaccination studies. *J Leukoc Biol* 2008;84:1364–73. doi:10.1189/jlb.0208092.
- [22] Lundberg K, Albrekt A-S, Nelissen I, Santegoets S, de Gruijl TD, Gibbs S, et al. Transcriptional profiling of human dendritic cell populations and models--unique profiles of *in vitro* dendritic cells and implications on functionality and applicability. *PLoS One* 2013;8:e52875. doi:10.1371/journal.pone.0052875.
- [23] Hitzler M, Bergert A, Luch A, Peiser M. Evaluation of selected biomarkers for the detection of chemical sensitization in human skin: A comparative study applying THP-1, MUTZ-3 and primary dendritic cells in culture. *Toxicol Vitr* 2013;27:1659–69. doi:10.1016/j.tiv.2013.04.009.
- [24] Brodin P, Davis MM. Human immune system variation. *Nat Rev Immunol* 2017;17:21–9. doi:10.1038/nri.2016.125.
- [25] Hynds RE, Vladimirov E, Janes SM. The secret lives of cancer cell lines. *Dis Model Mech* 2018;11:dmm037366. doi:10.1242/dmm.037366.
- [26] Stoel M, Pool J, de Vries-Idema J, Zaaraoui-Boutahar F, Bijl M, Andeweg AC, et al. Innate Responses Induced by Whole Inactivated Virus or Subunit Influenza Vaccines in Cultured Dendritic Cells Correlate with Immune Responses *In vivo*. *PLoS One* 2015;10:e0125228.
- [27] Nicholson KG, Tyrrell DAJ, Harrison P, Potter CW, Jennings R, Clark A, et al. Clinical studies of monovalent inactivated whole virus and subunit A/USSR/77 (H1N1) vaccine: serological responses and clinical reactions. *J Biol Stand* 1979;7:123–36. doi:10.1016/S0092-1157(79)80044-X.
- [28] Stephenson I, Nicholson KG, Glöck R, Mischler R, Newman RW, Palache AM, et al. Safety and antigenicity of whole virus and subunit influenza A/Hong Kong/1073/99 (H9N2) vaccine in healthy adults: phase I randomised trial. *The Lancet* 2003;362:1959–66.
- [29] Netea MG, Balkwill F, Chonchol M, Cominelli F, Donath MY, Giamarellos-Bourboulis EJ, et al. A guiding map for inflammation. *Nat Immunol* 2017;18:826–31. doi:10.1038/ni.3790.
- [30] Natoli G, Ostuni R. Adaptation and memory in immune responses. *Nat Immunol* 2019;20:783–92. doi:10.1038/s41590-019-0399-9.
- [31] Farber DL, Netea MG, Radbruch A, Rajewsky K, Zinkernagel RM. Immunological memory: lessons from the past and a look to the future. *Nat Rev Immunol* 2016;16:124–8. doi:10.1038/nri.2016.13.
- [32] Grødeland G, Fossum E, Bogen B. Polarizing T and B Cell Responses by APC-Targeted Subunit Vaccines. *Front Immunol* 2015;6:367. doi:10.3389/fimmu.2015.00367.
- [33] Clem AS. Fundamentals of Vaccine Immunology. *J Glob Infect Dis* 2011;3:73. doi:10.4103/0974-777X.77299.
- [34] Rosendahl Huber S, van Beek J, de Jonge J, Luytjes W, van Baarle D. T Cell Responses to Viral Infections – Opportunities for Peptide Vaccination. *Front Immunol* 2014;5:171. doi:10.3389/fimmu.2014.00171.
- [35] Halbroth BR, Heil A, Distler E, Dass M, Wagner EM, Plachter B, et al. Superior *In vitro* Stimulation of Human CD8⁺ T-Cells by Whole Virus versus Split Virus Influenza Vaccines. *PLoS One* 2014;9:e103392. doi:10.1371/journal.pone.0103392.
- [36] Reyes-Sandoval A, Pearson F, Todryk S, Ewe K. Potency assays for novel T-cell-inducing vaccines against malaria. *Curr Opin Mol Ther* 2009;11:72–80.
- [37] Clemens E, van de Sandt C, Wong S, Wakim L, Valkenburg S. Harnessing the Power of T Cells: The Promising Hope for a Universal Influenza Vaccine. *Vaccines* 2018;6:18. doi:10.3390/vaccines6020018.
- [38] Demers KR, Makedonas G, Buggert M, Eller MA, Ratcliffe SJ, Goonetilleke N, et al. Temporal Dynamics of CD8⁺ T Cell Effector Responses during Primary HIV Infection. *PLOS Pathog* 2016;12:e1005805.
- [39] Lin P-H, Wong W-I, Wang Y-L, Hsieh M-P, Lu C, Liang C-Y, et al. Vaccine-induced antigen-specific regulatory T cells attenuate the antiviral immunity against acute influenza virus infection. *Mucosal Immunol* 2018. doi:10.1038/s41385-018-0004-9.
- [40] Soema PC, van Riet E, Kersten G, Amorij J-P. Development of Cross-Protective Influenza A Vaccines Based on Cellular Responses. *Front Immunol* 2015;6:237.
- [41] Soema PC, van Riet E, Kersten G, Amorij

- J-P. Development of Cross-Protective Influenza A Vaccines Based on Cellular Responses. *Front Immunol* 2015;6:237. doi:10.3389/fimmu.2015.00237.
- [42] Gilbert SC. T-cell-inducing vaccines - what's the future. *Immunology* 2012;135:19-26. doi:10.1111/j.1365-2567.2011.03517.x.
- [43] He J, Tsai LM, Leong YA, Hu X, Ma CS, Chevalier N, et al. Circulating Precursor CCR7loPD-1hi CXCR5⁺ CD4⁺ T Cells Indicate Tfh Cell Activity and Promote Antibody Responses upon Antigen Reexposure. *Immunity* 2013;39:770-81. doi:10.1016/J.IMMUNI.2013.09.007.
- [44] Schmitt N, Ueno H. Blood Tfh Cells Come with Colors. *Immunity* 2013;39:629-30. doi:10.1016/J.IMMUNI.2013.09.011.
- [45] Bentebibel S-E, Khurana S, Schmitt N, Kurup P, Mueller C, Obermoser G, et al. ICOS⁺PD-1⁺CXCR3⁺ T Follicular Helper Cells Contribute To The Generation Of High-Avidity Antibodies Following Influenza Vaccination. *Sci Rep* 2016;6:26494. doi:10.1038/srep26494.
- [46] Bentebibel S-E, Lopez S, Obermoser G, Schmitt N, Mueller C, Harrod C, et al. Induction Of ICOS⁺CXCR3⁺CXCR5⁺ TH Cells Correlates With Antibody Responses To Influenza Vaccination. *Sci Transl Med* 2013;5:176ra32 LP-176ra32. doi:10.1126/scitranslmed.3005191.
- [47] Komanduri K V, Thomas M, St. John LS, Xing D, Decker W, Robinson S, et al. Characterization of optimal T Cell/Dendritic Cell (DC) Co-Culture Conditions for *Ex vivo* Expansion of Antigen-Specific Human T Cells. *Blood* 2006;108:3654 LP - 3654. doi:https://doi.org/.
- [48] Höpken UE, Lehmann I, Droese J, Lipp M, Schüler T, Rehm A. The Ratio Between Dendritic Cells And T Cells Determines The Outcome Of Their Encounter: Proliferation Versus Deletion. *Eur J Immunol* 2005;35:2851-63. doi:10.1002/eji.200526298.
- [49] Moser JM, Sassano ER, Leistritz DC, Eatrides JM, Phogat S, Koff W, et al. Optimization Of A Dendritic Cell-Based Assay For The *In vitro* Priming Of Naïve Human CD4⁺ T Cells. *J Immunol Methods* 2010;353:8-19. doi:10.1016/j.jim.2009.11.006.
- [50] Yao-Chen L, Ling-Hong K, Bi-Zhen C, Kang-Sheng L. Construction of Influenza Virus siRNA Expression Vectors and Their Inhibitory Effects on Multiplication of Influenza Virus. *Avian Dis* 2005;49:562-73. doi:10.1637/7365-041205R2.1.
- [51] Schultz HS, Reedtz-Runge SL, Bäckström BT, Lamberth K, Pedersen CR, Kvarnhammar AM, et al. Quantitative Analysis Of The CD4⁺ T Cell Response To Therapeutic Antibodies In Healthy Donors Using A Novel T Cell:PBMC Assay. *PLoS One* 2017;12:e0178544. doi:10.1371/journal.pone.0178544.
- [52] Hillaire MLB, van Trierum SE, Bodewes R, van Baalen CA, van Binnendijk RS, Koopmans MP, et al. Characterization Of The Human CD8⁺ T Cell Response Following Infection With 2009 Pandemic Influenza H1N1 Virus. *J Virol* 2011;85:12057 LP - 12061. doi:10.1128/JVI.05204-11.
- [53] Sridhar S, Begom S, Bermingham A, Hoschler K, Adamson W, Carman W, et al. Cellular Immune Correlates Of Protection Against Symptomatic Pandemic Influenza. *Nat Med* 2013;19:1305. doi:10.1038/nm.3350.
- [54] Wherry EJ, Teichgräber V, Becker TC, Masopust D, Kaech SM, Antia R, et al. Lineage Relationship And Protective Immunity Of Memory CD8 T Cell Subsets. *Nat Immunol* 2003;4:225. doi:10.1038/ni889.
- [55] Xie D, McElhaney JE. Lower GrB⁺ CD62L^{high} CD8 T_{CM} effector lymphocyte response to influenza virus in older adults is associated with increased CD28^{null} CD8 T lymphocytes. *Mech Ageing Dev* 2007;128:392-400. doi:10.1016/J.MAD.2007.05.001.
- [56] Saurwein-Teissl M, Lung TL, Marx F, Gschösser C, Asch E, Blasko I, et al. Lack Of Antibody Production Following Immunization In Old Age: Association With CD8⁽⁺⁾CD28⁽⁻⁾ T Cell Clonal Expansions And An Imbalance In The Production Of T_{H1} And T_{H2} Cytokines. *J Immunol* 2002;168:5893 LP - 5899. doi:10.4049/jimmunol.168.11.5893.
- [57] Moris P, van der Most R, Leroux-Roels I, Clement F, Dramé M, Hanon E, et al. H5N1 Influenza Vaccine Formulated with AS03A Induces Strong Cross-Reactive and Polyfunctional CD4 T-Cell Responses. *J Clin Immunol* 2011;31:443-54. doi:10.1007/s10875-010-9490-6.
- [58] Budimir N, Haan A de, Meijerhof T, Waijer S, Boon L, Gostick E, et al. Critical Role of TLR7 Signaling in the Priming of Cross-Protective Cytotoxic T Lymphocyte Responses by a Whole Inactivated Influenza Virus Vaccine. *PLoS One* 2013;8. doi:10.1371/journal.pone.0063163.
- [59] Stephenson I, Nicholson KG, Glück R, Mischler R, Newman RW, Palache AM, et al. Safety and antigenicity of whole virus and subunit influenza A/Hong Kong/1073/99



- (H9N2) vaccine in healthy adults: phase I randomised trial. *Lancet* 2003;362:1959–66. doi:10.1016/S0140-6736(03)15014-3.
- [60] Treanor JJ, Campbell JD, Zangwill KM, Rowe T, Wolff M. Safety and Immunogenicity of an Inactivated Subvirion Influenza A (H5N1) Vaccine. *N Engl J Med* 2006;354:1343–51. doi:10.1056/NEJMoa055778.
- [61] Brady RC, Treanor JJ, Atmar RL, Keitel WA, Edelman R, Chen WH, et al. Safety and immunogenicity of a subvirion inactivated influenza A/H5N1 vaccine with or without aluminum hydroxide among healthy elderly adults. *Vaccine* 2009;27:5091–5. doi:10.1016/J.VACCINE.2009.06.057.
- [62] Belshe RB, Frey SE, Graham I, Mulligan MJ, Edupuganti S, Jackson LA, et al. Safety and Immunogenicity of Influenza A H5 Subunit Vaccines: Effect of Vaccine Schedule and Antigenic Variant. *J Infect Dis* 2011;203:666–73. doi:10.1093/infdis/jiq093.
- [63] Lin J, Zhang J, Dong X, Fang H, Chen J, Su N, et al. Safety and immunogenicity of an inactivated adjuvanted whole-virion influenza A (H5N1) vaccine: a phase I randomised controlled trial. *The Lancet* 2006;368:991–7.
- [64] Leroux-Roels I, Borkowski A, Vanwolleghem T, Dramé M, Clement F, Hons E, et al. Antigen sparing and cross-reactive immunity with an adjuvanted rH5N1 prototype pandemic influenza vaccine: a randomised controlled trial. *Lancet* 2007;370:580–9. doi:10.1016/S0140-6736(07)61297-5.
- [65] Phan TL, Ho VT, Vu MH, Nguyen TN, Duong HT, Holt R, et al. Clinical testing of an inactivated influenza A/H5N1 vaccine candidate in a double-blinded, placebo-controlled, randomized trial in healthy adults in Vietnam. *Vaccine* 2016;34:5449–56. doi:10.1016/j.vaccine.2016.08.055.
- [66] Couch RB, Decker WK, Utama B, Atmar RL, Niño D, Feng JQ, et al. Evaluations for *In vitro* Correlates of Immunogenicity of Inactivated Influenza A H5, H7 and H9 Vaccines in Humans. *PLoS One* 2012;7:e50830.
- [67] Morefield GL. A Rational, Systematic Approach for the Development of Vaccine Formulations. *AAPS J* 2011;13:191–200. doi:10.1208/s12248-011-9261-1.
- [68] Foged C. Subunit vaccines of the future: the need for safe, customized and optimized particulate delivery systems. *Ther Deliv* 2011;2:1057–77. doi:10.4155/tde.11.68.
- [69] O'Hagan DT, De Gregorio E. The path to a successful vaccine adjuvant – 'The long and winding road.' *Drug Discov Today* 2009;14:541–51. doi:10.1016/j.drudis.2009.02.009.
- [70] STORNI T, KUNDIG T, SENTI G, JOHANSEN P. Immunity in response to particulate antigen-delivery systems. *Adv Drug Deliv Rev* 2005;57:333–55. doi:10.1016/j.addr.2004.09.008.
- [71] Trovato M, De Berardinis P. Novel antigen delivery systems. *World J Virol* 2015;4:156–68. doi:10.5501/wjv.v4.i3.156.
- [72] O'Hagan DT, Valiante NM. Recent advances in the discovery and delivery of vaccine adjuvants. *Nat Rev Drug Discov* 2003;2:727–35. doi:10.1038/nrd1176.
- [73] Bachmann MF, Jennings GT. Vaccine delivery: a matter of size, geometry, kinetics and molecular patterns. *Nat Rev Immunol* 2010;10:787.
- [74] Herzog C, Hartmann K, Künzi V, Kürsteiner O, Mischler R, Lazar H, et al. Eleven years of Inflexal® V—a virosomal adjuvanted influenza vaccine. *Vaccine* 2009;27:4381–7. doi:https://doi.org/10.1016/j.vaccine.2009.05.029.
- [75] Van Damme P, Dionne M, Leroux-Roels G, Van Der Meeren O, Di Paolo E, Salaun B, et al. Persistence of HBsAg-specific antibodies and immune memory two to three decades after hepatitis B vaccination in adults. *J Viral Hepat* n.d.;0. doi:10.1111/jvh.13125.
- [76] Vartak A, Sucheck SJ. Recent Advances in Subunit Vaccine Carriers. *Vaccines* 2016;4. doi:10.3390/vaccines4020012.
- [77] Medaglini D, De Azero MR, Leroy O, Bietrix F, Denoel P. Innovation Partnership for a Roadmap on Vaccines in Europe (IPROVE): A vision for the vaccines of tomorrow. *Vaccine* 2018;36:1136–45. doi:10.1016/j.vaccine.2017.11.069.
- [78] Green S, Rothman A. Immunopathological mechanisms in dengue and dengue hemorrhagic fever. *Curr Opin Infect Dis* 2006;19:429–36. doi:10.1097/01.qco.0000244047.31135.fa.
- [79] Fletcher HA, Chatterjee M, Cooper A, Hussell T, Kaye PM, Prior J, et al. VALIDATE: Exploiting the synergy between complex intracellular pathogens to expedite vaccine research and development for tuberculosis,



leishmaniasis, melioidosis and leprosy. *Fl000Research* 2018;7:485. doi:10.12688/fl000research.14386.1.

[80] Koff WC, Burton DR, Johnson PR, Walker BD, King CR, Nabel GJ, et al. Accelerating next-generation vaccine development for global disease prevention. *Science* 2013;340:1232910. doi:10.1126/science.1232910.

[81] Nederlandse Voedsel- en Warenautoriteit (NVWA). Jaaroverzicht dierproeven en proefdieren. 2015.

[82] De Mattia F, Chapsal J-M, Descamps J, Halder M, Jarrett N, Kross I, et al. The consistency approach for quality control of vaccines—A strategy to improve quality control and implement 3Rs. *Biologicals* 2011;39:59–65. doi:10.1016/j.biologicals.2010.12.001.

[83] De Mattia F, Hendriksen C, Buchheit KH, Chapsal JM, Halder M, Lambripts D, et al. The vaccines consistency approach project: an EPAA initiative. *Pharmeur Bio Sci Notes* 2015;2015:30–56.

[84] European Vaccine Initiative. VAC2VAC | European Vaccine Initiative n.d. <http://www.euvaccine.eu/portfolio/project-index/vac2vac> (accessed July 28, 2019).

[85] Hendriksen C, Arciniega JL, Bruckner L, Chevalier M, Coppens E, Descamps J, et al. The consistency approach for the quality control of vaccines. *Biologicals* 2008;36:73–7. doi:10.1016/j.biologicals.2007.05.002.

[86] Geeraedts F, Bungener L, Pool J, W TV, Wilschut J, Huckriede A. Whole inactivated virus influenza vaccine is superior to subunit vaccine in inducing immune responses and secretion of proinflammatory cytokines by DCs. *Influ Other Respi Viruses* 2008;2:41–51. doi:10.1111/j.1750-2659.2008.00038.x.

[87] Geeraedts F, Goutagny N, Hornung V, Severa M, de Haan A, Pool J, et al. Superior Immunogenicity of Inactivated Whole Virus H5N1 Influenza Vaccine is Primarily Controlled by Toll-like Receptor Signalling. *PLoS Pathog* 2008;4:e1000138. doi:10.1371/journal.ppat.1000138.

[88] Shi W, Liao Y, Willis SN, Taubenheim N, Inouye M, Tarlinton DM, et al. Transcriptional profiling of mouse B cell terminal differentiation defines a signature for antibody-secreting plasma cells. *Nat Immunol* 2015;16:663–73. doi:10.1038/ni.3154.

[89] Nakaya HI, Clutterbuck E, Kazmin D, Wang L, Cortese M, Bosinger SE, et al. Systems biology of immunity to MF59-adjuvanted

versus nonadjuvanted trivalent seasonal influenza vaccines in early childhood. *Proc Natl Acad Sci U S A* 2016;113:1853–8. doi:10.1073/pnas.1519690113.

[90] Nakaya HI, Wrammert J, Lee EK, Racioppi L, Marie-Kunze S, Haining WN, et al. Systems biology of vaccination for seasonal influenza in humans. *Nat Immunol* 2011;12:786–95. doi:10.1038/ni.2067.

[91] Tan Y, Tamayo P, Nakaya H, Pulendran B, Mesirov JP, Haining WN. Gene signatures related to B-cell proliferation predict influenza vaccine-induced antibody response. *Eur J Immunol* 2014;44:285–95. doi:10.1002/eji.201343657.

[92] Nakaya HI, Wrammert J, Lee EK, Racioppi L, Marie-Kunze S, Haining WN, et al. Systems biology of vaccination for seasonal influenza in humans. *Nat Immunol* 2011;12:786–95. doi:10.1038/ni.2067.

[93] Querec T, Bennouna S, Alkan S, Laouar Y, Gorden K, Flavell R, et al. Yellow fever vaccine YF-17D activates multiple dendritic cell subsets via TLR2, 7, 8, and 9 to stimulate polyvalent immunity. *J Exp Med* 2006;203:413–24. doi:10.1084/jem.20051720.

[94] Querec TD, Akondy RS, Lee EK, Cao W, Nakaya HI, Teuwen D, et al. Systems biology approach predicts immunogenicity of the yellow fever vaccine in humans. *Nat Immunol* 2009;10:116–25. doi:10.1038/ni.1688.

[95] Anderson J, Olafsdottir TA, Kratochvil S, McKay PF, Östensson M, Persson J, et al. Molecular Signatures of a TLR4 Agonist-Adjuvanted HIV-1 Vaccine Candidate in Humans. *Front Immunol* 2018;9:301. doi:10.3389/fimmu.2018.00301.

[96] Strouts FR, Popper SJ, Partidos CD, Stinchcomb DT, Osorio JE, Relman DA. Early Transcriptional Signatures of the Immune Response to a Live Attenuated Tetravalent Dengue Vaccine Candidate in Non-human Primates. *PLoS Negl Trop Dis* 2016;10:e0004731. doi:10.1371/journal.pntd.0004731.

[97] Reichtien A, Richert L, Lorenzo H, Martrus G, Hejblum B, Dahlke C, et al. Systems Vaccinology Identifies an Early Innate Immune Signature as a Correlate of Antibody Responses to the Ebola Vaccine rVSV-ZEBOV. *Cell Rep* 2017;20:2251–61. doi:10.1016/j.celrep.2017.08.023.

[98] Ravindran R, Khan N, Nakaya HI, Li S, Loebbermann J, Maddur MS, et al. Vaccine activation of the nutrient sensor



- GCN2 in dendritic cells enhances antigen presentation. *Science* 2014;343:313–7. doi:10.1126/science.1246829.
- [99] Nakaya H, Wrammert J, Lee E, Racioppi L, Marie-Kunze S, Haining W, et al. Systems Biology of Seasonal Influenza Vaccination in Humans. *Nat Immunol* 2011;12:786–95.
- [100] Spensieri F, Borgogni E, Zedda L, Bardelli M, Buricchi F, Volpini G, et al. Human Circulating Influenza-CD4⁺ ICOS⁺IL-21⁺ T Cells Expand After Vaccination, Exert Helper Function, And Predict Antibody Responses. *Proc Natl Acad Sci* 2013;110:14330–5. doi:10.1073/PNAS.1311998110.
- [101] Abbott RK, Lee JH, Menis S, Skog P, Rossi M, Ota T, et al. Precursor Frequency and Affinity Determine B Cell Competitive Fitness in Germinal Centers, Tested with Germ-line-Targeting HIV Vaccine Immunogens. *Immunity* 2018;48:133–146.e6. doi:10.1016/j.immuni.2017.11.023.
- [102] Pallikkuth S, de Armas LR, Rinaldi S, George VK, Pan L, Arheart KL, et al. Dysfunctional peripheral T follicular helper cells dominate in people with impaired influenza vaccine responses: Results from the FLORAH study. *PLOS Biol* 2019;17:e3000257. doi:10.1371/journal.pbio.3000257.
- [103] Cubas RA, Mudd JC, Savoye A-L, Perreau M, van Grevenynghe J, Metcalf T, et al. Inadequate T follicular cell help impairs B cell immunity during HIV infection. *Nat Med* 2013;19:494–9. doi:10.1038/nm.3109.
- [104] Perreau M, Savoye A-L, De Crignis E, Corpataux J-M, Cubas R, Haddad EK, et al. Follicular helper T cells serve as the major CD4 T cell compartment for HIV-1 infection, replication, and production. *J Exp Med* 2013;210:143–56. doi:10.1084/jem.20121932.
- [105] Ladhani S, Slack MPE, Ramsay ME, Heath PT, McIntyre PB, Diez-Domingo J, et al. Haemophilus influenzae serotype b conjugate vaccine failure in twelve countries with established national childhood immunization programmes. *Clin Microbiol Infect* 2010;16:948–54. doi:10.1111/j.1469-0691.2009.02945.x.
- [106] McLean HQ, Chow BDW, VanWormer JJ, King JP, Belongia EA. Effect of Statin Use on Influenza Vaccine Effectiveness. *J Infect Dis* 2016;214:1150–8. doi:10.1093/infdis/jiw335.
- [107] Long JE, Drayson MT, Taylor AE, Toellner KM, Lord JM, Phillips AC. Morning vaccination enhances antibody response over afternoon vaccination: A cluster-randomised trial. *Vaccine* 2016;34:2679–85. doi:10.1016/j.vaccine.2016.04.032.
- [108] De Serres G, Boulianne N, Defay F, Brousseau N, Benoît M, Lacoursière S, et al. Higher Risk of Measles When the First Dose of a 2-Dose Schedule of Measles Vaccine Is Given at 12–14 Months Versus 15 Months of Age. *Clin Infect Dis* 2012;55:394–402. doi:10.1093/cid/cis439.
- [109] Trotter CL, Andrews NJ, Kaczmarski EB, Miller E, Ramsay ME. Effectiveness of meningococcal serogroup C conjugate vaccine 4 years after introduction. *Lancet* 2004;364:365–7. doi:10.1016/S0140-6736(04)16725-1.
- [110] McElhane JE, Kuchel GA, Zhou X, Swain SL, Haynes L. T-Cell Immunity to Influenza in Older Adults: A Pathophysiological Framework for Development of More Effective Vaccines. *Front Immunol* 2016;7:41. doi:10.3389/fimmu.2016.00041.
- [111] WHO. WHO | Vaccines and diseases. WHO 2019. <https://www.who.int/immunization/diseases/en/> (accessed July 21, 2019).
- [112] Ventura MT, Casciaro M, Gangemi S, Buquicchio R. Immunosenescence in aging: between immune cells depletion and cytokines up-regulation. *Clin Mol Allergy* 2017;15:21. doi:10.1186/s12948-017-0077-0.
- [113] Nikolich-Zugich J. The twilight of immunity: emerging concepts in aging of the immune system. *Nat Immunol* 2018;19:10–9. doi:10.1038/s41590-017-0006-x.
- [114] Goodwin K, Viboud C, Simonsen L. Antibody response to influenza vaccination in the elderly: A quantitative review. *Vaccine* 2006;24:1159–69. doi:10.1016/J.VACCINE.2005.08.105.
- [115] Trzonkowski P, Myśliwska J, Szmit E, Wieckiewicz J, Lukaszuk K, Brydak LB, et al. Association between cytomegalovirus infection, enhanced proinflammatory response and low level of anti-hemagglutinins during the anti-influenza vaccination--an impact of immunosenescence. *Vaccine* 2003;21:3826–36.
- [116] Lynch HE, Goldberg GL, Chidgey A, Van den Brink MRM, Boyd R, Sempowski GD. Thymic involution and immune reconstitution. *Trends Immunol* 2009;30:366–73. doi:10.1016/j.it.2009.04.003.
- [117] Shanley DP, Aw D, Manley NR, Palmer DB. An evolutionary perspective on the mechanisms of immunosenescence. *Trends*



- Immunol 2009;30:374–81. doi:10.1016/j.it.2009.05.001.
- [118] Hearps AC, Martin GE, Angelovich TA, Cheng W-J, Maisa A, Landay AL, et al. Aging is associated with chronic innate immune activation and dysregulation of monocyte phenotype and function. *Aging Cell* 2012;11:867–75. doi:10.1111/j.1474-9726.2012.00851.x.
- [119] Thompson KM, Tebbens RJD. Retrospective Cost-Effectiveness Analyses for Polio Vaccination in the United States. *Risk Anal* 2006;26:1423–40. doi:10.1111/j.1539-6924.2006.00831.x.
- [120] Kong K-F, Delroux K, Wang X, Qian F, Arjona A, Malawista SE, et al. Dysregulation of TLR3 impairs the innate immune response to West Nile virus in the elderly. *J Virol* 2008;82:7613–23. doi:10.1128/JVI.00618-08.
- [121] McLaughlin JM, McGinnis JJ, Tan L, Mercatante A, Fortuna J. Estimated Human and Economic Burden of Four Major Adult Vaccine-Preventable Diseases in the United States, 2013. *J Prim Prev* 2015;36:259–73. doi:10.1007/s10935-015-0394-3.
- [122] Herpes Zoster and Functional Decline Consortium. Functional decline and herpes zoster in older people: an interplay of multiple factors. *Aging Clin Exp Res* 2015;27:757–65. doi:10.1007/s40520-015-0450-0.
- [123] Janssens J-P. Pneumonia in the elderly (geriatric) population. *Curr Opin Pulm Med* 2005;11:226–30.
- [124] Lorenzo EC, Bartley JM, Haynes L. The impact of aging on CD4⁺ T cell responses to influenza infection. *Biogerontology* 2018;19:437–46. doi:10.1007/s10522-018-9754-8.
- [125] Weinberger B, Grubeck-Loebenstein B. Vaccines for the elderly. *Clin Microbiol Infect* 2012;18:100–8. doi:10.1111/J.1469-0691.2012.03944.X.
- [126] Weinberger B. Vaccines for the elderly: current use and future challenges. *Immun Ageing* 2018;15:3. doi:10.1186/s12979-017-0107-2.
- [127] Chen WH, Kozlovsky BF, Effros RB, Grubeck-Loebenstein B, Edelman R, Szein MB. Vaccination in the elderly: an immunological perspective. *Trends Immunol* 2009;30:351. doi:10.1016/J.IT.2009.05.002.
- [128] Aspinall R, Del Giudice G, Effros RB, Grubeck-Loebenstein B, Sambhara S. Challenges for vaccination in the elderly. *Immun Ageing* 2007;4:9. doi:10.1186/1742-4933-4-9.
- [129] Beyer WEP, McElhaney J, Smith DJ, Monto AS, Nguyen-Van-Tam JS, Osterhaus ADME. Cochrane re-arranged: Support for policies to vaccinate elderly people against influenza. *Vaccine* 2013;31:6030–3. doi:10.1016/j.vaccine.2013.09.063.
- [130] Osterholm MT, Kelley NS, Sommer A, Belongia EA. Efficacy and effectiveness of influenza vaccines: a systematic review and meta-analysis. *Lancet Infect Dis* 2012;12:36–44. doi:10.1016/S1473-3099(11)70295-X.
- [131] Belongia EA, Simpson MD, King JP, Sundaram ME, Kelley NS, Osterholm MT, et al. Variable influenza vaccine effectiveness by subtype: a systematic review and meta-analysis of test-negative design studies. *Lancet Infect Dis* 2016;16:942–51. doi:10.1016/S1473-3099(16)00129-8.
- [132] Gostic KM, Ambrose M, Worobey M, Lloyd-Smith JO. Potent protection against H5N1 and H7N9 influenza via childhood hemagglutinin imprinting. *Science* (80-) 2016;354:722–6. doi:10.1126/science.aag1322.
- [133] Weisel F, Shlomchik M. Memory B Cells of Mice and Humans. *Annu Rev Immunol* 2017;35:255–84. doi:10.1146/annurev-immunol-041015-055531.
- [134] Hong S, Zhang Z, Liu H, Tian M, Zhu X, Zhang Z, et al. B Cells Are the Dominant Antigen-Presenting Cells that Activate Naive CD4⁺ T Cells upon Immunization with a Virus-Derived Nanoparticle Antigen. *Immunity* 2018;49:695–708.e4. doi:10.1016/j.immuni.2018.08.012.
- [135] Rossetti RAM, Lorenzi NPC, Yokochi K, Rosa MBS de F, Benevides L, Margarido PFR, et al. B lymphocytes can be activated to act as antigen presenting cells to promote anti-tumor responses. *PLoS One* 2018;13:e0199034. doi:10.1371/journal.pone.0199034.
- [136] Chen X, Jensen PE. The role of B lymphocytes as antigen-presenting cells. *Arch Immunol Ther Exp (Warsz)* 2008;56:77–83. doi:10.1007/s00005-008-0014-5.
- [137] Popi AF, Longo-Maugéri IM, Mariano M. An Overview of B-1 Cells as Antigen-Presenting Cells. *Front Immunol* 2016;7:138. doi:10.3389/fimmu.2016.00138.
- [138] Pasare C, Medzhitov R. Control of B-cell responses by Toll-like receptors. *Nature* 2005;438:364–8. doi:10.1038/nature04267.
- [139] Ruprecht CR, Lanzavecchia A. Toll-like receptor stimulation as a third signal



required for activation of human naive B cells. *Eur J Immunol* 2006;36:810–6. doi:10.1002/eji.200535744.

[140] Cognasse F, Hamzeh-Cognasse H, Lafarge S, Chavarin P, Pozzetto B, Richard Y, et al. Identification of two subpopulations of purified human blood B cells, CD27⁺ CD23⁺ and CD27^{high} CD80⁺, that strongly express cell surface Toll-like receptor 9 and secrete high levels of interleukin-6. *Immunology* 2008;125:430–7. doi:10.1111/j.1365-2567.2008.02844.x.

[141] Bernasconi NL, Onai N, Lanzavecchia A. A role for Toll-like receptors in acquired immunity: up-regulation of TLR9 by BCR triggering in naive B cells and constitutive expression in memory B cells. *Blood* 2003;101:4500 LP – 4504.

[142] Jung S, Unutmaz D, Wong P, Sano G-I, De los Santos K, Sparwasser T, et al. *In vivo* Depletion of CD11c⁺ Dendritic Cells Abrogates Priming of CD8⁺ T Cells by Exogenous Cell-Associated Antigens. *Immunity* 2002;17:211–20. doi:10.1016/S1074-7613(02)00365-5.

[143] Bernasconi NL, Traggiai E, Lanzavecchia A. Maintenance of Serological Memory by Polyclonal Activation of Human Memory B Cells. *Science* (80-) 2002;298:2199–202. doi:10.1126/science.1076071.

[144] Pone EJ, Zhang J, Mai T, White CA, Li G, Sakakura JK, et al. BCR-signalling synergizes with TLR-signalling for induction of AID and immunoglobulin class-switching through the non-canonical NF- κ B pathway. *Nat Commun* 2012;3:767. doi:10.1038/ncomms1769.

[145] Aranburu A, Ceccarelli S, Giorda E, Lasorella R, Ballatore G, Carsetti R. TLR Ligation Triggers Somatic Hypermutation in Transitional B Cells Inducing the Generation of IgM Memory B Cells. *J Immunol* 2010;178:3293–301. doi:10.4049/jimmunol.1002722.

[146] Ueda Y, Liao D, Yang K, Patel A, Kelsoe G. T-independent activation-induced cytidine deaminase expression, class-switch recombination, and antibody production by immature/transitional T B cells. *J Immunol* 2007;178:3593–601. doi:10.4049/jimmunol.178.6.3593.

[147] Lane P, Brouck T, Hubele S, Padovan E, Lanzavecchia A, McConnell F. Soluble CD40 ligand can replace the normal T cell-derived CD40 ligand signal to B cells in T cell-dependent activation. *J Exp Med* 1993;177:1209–13. doi:10.1084/jem.177.4.1209.

[148] Sasaki S, Jaimes MC, Holmes TH, Dekker CL, Mahmood K, Kemble GW, et al. Comparison of the Influenza Virus-Specific Effector and Memory B-Cell Responses to Immunization of Children and Adults with Live Attenuated or Inactivated Influenza Virus Vaccines. *J Virol* 2007;81:215–28. doi:10.1128/JVI.01957-06.

[149] Li G-M, Chiu C, Wrammert J, McCausland M, Andrews SF, Zheng N-Y, et al. Pandemic H1N1 influenza vaccine induces a recall response in humans that favors broadly cross-reactive memory B cells. *Proc Natl Acad Sci U S A* 2012;109:9047–52. doi:10.1073/pnas.1118979109.

[150] Hirota Y, Kaji M, Ide S, Kajiwarra J, Kataoka K, Goto S, et al. Antibody efficacy as a keen index to evaluate influenza vaccine effectiveness. *Vaccine* 1997;15:962–7.

[151] Bernasconi NL, Onai N, Lanzavecchia A. A role for Toll-like receptors in acquired immunity: up-regulation of TLR9 by BCR triggering in naive B cells and constitutive expression in memory B cells. *Blood* 2003;101:4500–4. doi:10.1182/blood-2002-11-3569.

[152] Preite S, Baumjohann D, Foglierini M, Basso C, Ronchi F, Rodriguez BMF, et al. Somatic mutations and affinity maturation are impaired by excessive numbers of T follicular helper cells and restored by Treg cells or memory T cells. *Eur J Immunol* 2015;45:3010–21. doi:10.1002/eji.201545920.

[153] Geffroy-Luseau A, Chiron D, Descamps G, Jégo G, Amiot M, Pellat-Deceunynck C. TLR9 Ligand Induces the Generation of CD20⁺ Plasmablasts and Plasma Cells from CD27⁺ Memory B-Cells. *Front Immunol* 2011;2:83.

[154] Jung J, Yi A-K, Zhang X, Choe J, Li L, Choi YS. Distinct Response of Human B Cell Subpopulations in Recognition of an Innate Immune Signal, CpG DNA. *J Immunol* 2002;169:2368–73. doi:10.4049/jimmunol.169.5.2368.

[155] Glaum MC, Narula S, Song D, Zheng Y, Anderson AL, Pletcher CH, et al. Toll-like receptor 7-induced naive human B-cell differentiation and immunoglobulin production. *J Allergy Clin Immunol* 2009;123:224–230.e4. doi:10.1016/j.jaci.2008.09.018.

[156] Borrebaeck CA, Danielsson L, Moller SA. Human monoclonal antibodies produced by primary *in vitro* immunization of peripheral blood lymphocytes. *Proc*



Natl Acad Sci 1988;85:3995–9. doi:10.1073/pnas.85.11.3995.

[157] Eckl-Dorna J, Batista FD. BCR-mediated uptake of antigen linked to TLR9 ligand stimulates B-cell proliferation and antigen-specific plasma cell formation. *Blood* 2009;113:3969–77. doi:10.1182/blood-2008-10-185421.

[158] Sanjuan Nandin I, Fong C, Deantonio C, Torreno-Pina JA, Pecetta S, Maldonado P, et al. Novel *in vitro* booster vaccination to rapidly generate antigen-specific human monoclonal antibodies. *J Exp Med* 2017;214:2471–90. doi:10.1084/jem.20170633.

[159] Fecteau JF, Néron S. CD40 Stimulation of Human Peripheral B Lymphocytes: Distinct Response from Naive and Memory Cells. *J Immunol* 2003;171:4621–9. doi:10.4049/jimmunol.171.9.4621.

[160] Takemori T, Kaji T, Takahashi Y, Shimoda M, Rajewsky K. Generation of memory B cells inside and outside germinal centers. *Eur J Immunol* 2014;44:1258–64. doi:10.1002/eji.201343716.

[161] Schwickert TA, Victora GD, Fooksman DR, Kamphorst AO, Mugnier MR, Gitlin AD, et al. A dynamic T cell-limited checkpoint regulates affinity-dependent B cell entry into the germinal center. *J Exp Med* 2011;208:1243–52. doi:10.1084/jem.20102477.

[162] Querec T, Bennouna S, Alkan S, Laouar Y, Gorden K, Flavell R, et al. Yellow fever vaccine YF-17D activates multiple dendritic cell subsets via TLR2, 7, 8, and 9 to stimulate polyvalent immunity. *J Exp Med* 2006;203:413–24.

[163] Pulendran B, Li S, Nakaya HI. Systems vaccinology. *Immunity* 2010;33:516–29. doi:10.1016/j.immuni.2010.10.006.

[164] Kasturi S, Skountzou I, Albrecht R, Koutsouanos D, Hua T, Nakaya H, et al. Programming the magnitude and persistence of antibody responses with innate immunity. *Nature* 2011;470:543–47.

[165] Theiler M, Smith HH. The Use Of Yellow Fever Virus Modified By *In vitro* Cultivation For Human Immunization. *J Exp Med* 1937;65:787–800.

[166] Koutsakos M, Nguyen THO, Kedzierska K. With a Little Help from T Follicular Helper Friends: Humoral Immunity to Influenza Vaccination. *J Immunol* 2019;202:360–7. doi:10.4049/jimmunol.1800986.

[167] Bucasas KL, Franco LM, Shaw CA, Bray MS, Wells JM, Niño D, et al. Early patterns of gene expression correlate with the humoral immune response to influenza vaccination in humans. *J Infect Dis* 2011;203:921–9. doi:10.1093/infdis/jiq156.

[168] Obermoser G, Presnell S, Domico K, Xu H, Wang Y, Anguiano E, et al. Systems Scale Interactive Exploration Reveals Quantitative and Qualitative Differences in Response to Influenza and Pneumococcal Vaccines. *Immunity* 2013;38:831–44. doi:10.1016/j.immuni.2012.12.008.

[169] Spensieri F, Siena E, Borgogni E, Zedda L, Cantisani R, Chiappini N, et al. Early Rise of Blood T Follicular Helper Cell Subsets and Baseline Immunity as Predictors of Persisting Late Functional Antibody Responses to Vaccination in Humans. *PLoS One* 2016;11:e0157066. doi:10.1371/journal.pone.0157066.

[170] Dasgupta G, BenMohamed L. Of mice and not humans: How reliable are animal models for evaluation of herpes CD8+T cell-epitopes-based immunotherapeutic vaccine candidates? *Vaccine* 2011;29:5824–36. doi:10.1016/j.vaccine.2011.06.083.

[171] van der Worp HB, Howells DW, Sena ES, Porritt MJ, Rewell S, O'Collins V, et al. Can Animal Models of Disease Reliably Inform Human Studies? *PLoS Med* 2010;7:e1000245. doi:10.1371/journal.pmed.1000245.172] Barré-Sinoussi F, Montagnutelli X. Animal models are essential to biological research: issues and perspectives. *Futur Sci OA* 2015;1:fso.15.63. doi:10.4155/fso.15.63.



